

65777 – 16.5 grams
65778 – 12.2 grams
Poikilitic Impact Melt Breccia




Figure 1: Photo of 65777. Sample is 3.5 cm. S72-48813




Figure 2: Thin section photo of 65777 (from warner et al. 1976).

Introduction

65777 and 65778 were collected as a rake sample from station 5 – see section on 65701.

65777 has been dated at 3.7 b.y.




Figure 3: Pyroxene and olivine composition of 65777 (Warner et al. 1976).

Petrography

Keil et al. (1972) and Warner et al. (1976) found 65777 and 65778 were coherent crystalline impact melt breccias with poikilitic texture. The interlocking pigeonite oikocrysts include rounded grains of augite




Figure 4: Photo of thin section 65777,5 by C Meyer. 2 mm across



Figure 5: Photo of 65778. Scale is in mm.. S72-48815



Figure 6: Photomicrograph of thin section of 65778 (from Warner et al. 1976).

and olivine, and laths and relict clasts of plagioclase (figures 2, 4 and 6).

Reimold and Borchardt (1984) studied the petrology of 65777, comparing it with that of 65015 and 62235, which are similar rocks. They found 11 % of the sample was made of small areas of sub-ophitic basalt.




Figure 7: Composition of olivine and pyroxene in 65778 (Warner et al. 1976).

Compositional variation of Apollo 16 impact-melt rocks is discussed by Korotev (1994).

Chemistry

Warner et al. (1976) and Laul and Schmitt (1973) reported the composition of 65777 (table 1). It is enriched in trace elements characteristic of KREEP. Reimold and Borchardt (1984) determined the composition of regions within 65777 that had distinctive melt texture.

Radiogenic age dating

Schaeffer and Schaeffer (1977) determined an Ar/Ar age of 3.70 ± 0.02 b.y. for 65777 (figure 8). Norman et al. (2006) should have included these samples in their study.

Cosmogenic isotopes and exposure ages

Schaeffer and Schaeffer (1977) determined an exposure age of 8 m.y. for 65777.

Processing

There are 3 thin sections of 65777 and two for 65778.





Figure 8: Isochron diagram for K/Ar dating of 65777 (Schaeffer and Schaeffer (1977)).

References for 65777

Butler P. (1972a) Lunar Sample Information Catalog Apollo 16. Lunar Receiving Laboratory. MSC 03210 Curator's Catalog. pp. 370.

Dowty E., Green J.A., Hlava P.F., Keil K., Moore R.B., Nehru C.E., Prinz M. and Warner R.D. (1976) Electron microprobe analyses of minerals from Apollo 16 rake samples. Inst. Meteoritics Spec. Publ. No 14, 141 pp.. Univ. New Mex. ABQ

Keil K., Dowty E., Prinz M. and Bunch T.E. (1972) Description, classification and inventory of 151 Apollo 16 rake samples from the LM area and station 5. Curator's Catalog, JSC.

Korotev R.L. (1994) Compositional variation in Apollo 16 impact melt breccias and inferences for the geology and bombardment history of the central highlands of the Moon. *Geochim. Cosmochim. Acta* **58**, 3931-3969.

Laul J.C. and Schmitt R.A. (1973b) Chemical composition of Apollo 15, 16, and 17 samples. *Proc. 4th Lunar Sci. Conf.* 1349-1367.

LSPET (1973b) The Apollo 16 lunar samples: Petrographic and chemical description. *Science* **179**, 23-34.

Table 1. Chemical composition of 65777

reference	Laul73	Warner76	Reimold84 subophitic areas
weight			
SiO ₂ %		47.7	(b) 46.42
TiO ₂	1.2	(a) 1.21	(b) 2.67
Al ₂ O ₃	18.5	(a) 20.3	(b) 17.18
FeO	9	(a) 7.7	(b) 7.78
MnO	0.106	(a) 0.07	(b) 0.04
MgO	10	(a) 11	(b) 11.34
CaO	11.3	(a) 12.1	(b) 10.96
Na ₂ O	0.66	(a) 0.71	(b) 0.44
K ₂ O	0.37	(a) 0.45	(b) 1.76
P ₂ O ₅			0.43 (b) 1.38
S %			0.42 (b)
sum			
Sc ppm	14	(a)	
V	35	(a)	
Cr	1164	(a)	
Co	59	(a)	
Ni	1100	(a)	
Cu			
Zn			
Ga			
Ge ppb			
As			
Se			
Rb			
Sr			
Y			
Zr	620	(a) 1200	(b)
Nb			
Mo			
Ru			
Rh			
Pd ppb			
Ag ppb			
Cd ppb			
In ppb			
Sn ppb			
Sb ppb			
Te ppb			
Cs ppm			
Ba	530	(a)	
La	53	(a)	
Ce	145	(a)	
Pr			
Nd	90	(a)	
Sm	23	(a)	
Eu	2.03	(a)	
Gd			
Tb	4.5	(a)	
Dy	30	(a)	
Ho			
Er			
Tm			
Yb	15	(a)	
Lu	2.1	(a)	
Hf	17	(a)	
Ta	2	(a)	
W ppb			
Re ppb			
Os ppb			
Ir ppb	17	(a)	
Pt ppb			
Au ppb	22	(a)	
Th ppm	8.8	(a)	
U ppm	3	(a)	
technique:	(a) INAA, (b) broad beam e probe		

Table 2. Chemical composition of 65778.

reference	Warner76
weight	
SiO ₂ %	47.3 (a)
TiO ₂	0.88 (a)
Al ₂ O ₃	21.6 (a)
FeO	6.8 (a)
MnO	0.06 (a)
MgO	9.7 (a)
CaO	12.5 (a)
Na ₂ O	0.52 (a)
K ₂ O	0.29 (a)
P ₂ O ₅	0.27 (a)
S %	
sum	
(a) DBA	

LSPET (1972c) Preliminary examination of lunar samples.
In Apollo 16 Preliminary Science Report. NASA SP-315, 7-1—7-58.

Reimold W.U. and Borchardt R. (1984) Subophitic lithologies in KREEP-rich poikilitic impact melt rocks from Cayley Plains, Apollo 16 – remnants of a volcanic Highland curst? *Earth Planet. Sci. Lett.* **67**, 9-18.

Reimold W.U. and Reimold J.N. (1984) The mineralogical, chemical and chronological characteristics of the crystalline Apollo 16 impact melt rocks. *Fortschr. Mineral.* **62**, 269-301.

Reimold W.U., Nyquist L.E., Bansal B.M., Wooden J.L., Shih C.-Y., Wiesmann H. and Mackinnon I.D.R. (1985) Isotope analysis of crystalline impact-melt rocks from Apollo 16 stations 11 and 13. North Ray Crater. *Proc. 15th Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **90**, C431-C448.

Ryder G. and Norman M.D. (1980) Catalog of Apollo 16 rocks (3 vol.). Curator's Office pub. #52, JSC #16904

Schaeffer G.A. and Schaeffer O.A. (1977a) ³⁹Ar/⁴⁰Ar ages of lunar rocks. *Proc. 8th Lunar Sci. Conf.* 2253-2300.

Sutton R.L. (1981) Documentation of Apollo 16 samples. In *Geology of the Apollo 16 area, central lunar highlands.* (Ulrich et al.) U.S.G.S. Prof. Paper 1048.

Warner R.D., Dowty E., Prinz M., Conrad G.H., Nehru C.E. and Keil K. (1976c) Catalog of Apollo 16 rake samples from the LM area and station 5. Spec. Publ. #13, UNM Institute of Meteoritics, Albuquerque. 87 pp.